
Borna disease virus and the evidence for human pathogenicity: a systematic review

R.M. CHALMERS, D.Rh. THOMAS and R.L. SALMON

From the NPHS Communicable Disease Surveillance Centre, Cardiff, UK

Received 23 July 2004

Summary

Background: Borna disease is a neurological viral disease of veterinary importance in central Europe, although Borna Disease virus (BDV) has been reported to be present in animals in most continents. The hypothesis that BDV is associated with human illness is controversial. However, should even a small fraction of mental illness be attributable to infection with BDV, this would be an important finding, not least because illness in that sub-population would, theoretically, be preventable.

Methods: We systematically reviewed the evidence: that BDV infects humans; for the role of BDV in human neuropsychiatric illness; to assess the suitability of currently available laboratory methods for human epidemiological studies.

Results: We identified 75 documents published before the end of January 2000, describing 50 human studies for BDV. There were five case studies and 44 (sero)prevalence studies, in a variety of patient groups. Nineteen prevalence studies (43%) investigated seroprevalence, 11 (25%) investigated viral prevalence and 14 (32%) investigated both. Seroprevalence ranged from 0% to 48%, and prevalence of virus or viral footprints from 0% to 82%.

Discussion: Although agreed gold standard tests and evidence for test specificity are lacking, there is evidence that humans are exposed to the virus. Further epidemiological studies are required to establish whether there are associations with disease.

Introduction

Borna disease was first described in the early 19th century as an 'equine brain disease with signs of agitation'¹ and obtained its name following an epidemic in military horses in Borna, Saxony, in 1885. Since then, this naturally occurring meningoencephalitic disease has been described mainly in horses, but also in sheep, cattle and other domesticated and wild animals and in a variety of experimentally infected host species.² The aetiological agent, Borna disease virus (BDV), also causes clinically definable and reproducible behavioural abnormalities in rats and non-human primates, some of which resemble neuropsychiatric disorders

in humans. Viral tropism for the limbic system, including excitatory fields in the hippocampus, has been reported.^{3,4} These areas of the brain govern emotions important to survival, visceral responses and olfactory sensations, and processes involved in memory. The detection of BDV antibodies and viral RNA in peripheral blood mononuclear cells (PBMCs) from psychiatric patients has furthered the suggestion that BDV is linked to human neuropsychiatric illness.⁵

Routes of transmission are unproven, but thought to be via excreta or nasal, saliva and conjunctival secretions, either directly or indirectly through

Address correspondence to Dr D.Rh. Thomas, NPHS Communicable Disease Surveillance Centre, Abton House, Wedal Road, Cardiff CF14 3QX. email: daniel.thomas@nphs.wales.nhs.uk

© The Author 2005. Published by Oxford University Press on behalf of the Association of Physicians.
All rights reserved. For Permissions, please email: journals.permissions@oupjournals.org

contaminated food and water. BDV RNA has been detected in secretions from horses by reverse transcriptase polymerase chain reaction (RT-PCR), both in the presence and absence of clinical illness.^{6,7} The detection of viral markers in blood raises the possibility of blood-borne transmission. The virus itself is an enveloped, non-segmented negative-strand RNA virus, Order Mononegavirales, Family Bornaviridae (ICV Jerusalem). Several features of its molecular biology are remarkable, and some are relevant to questions regarding the detection and spread of the virus, and the epidemiology of disease. For example, the virus replicates at lower levels than many other viruses, producing low numbers of infectious particles, although cell to cell transmission of incomplete particles may occur.⁸ Sensitive methods are therefore required for virus detection and identification. Demonstration of BDV antigens in populations of PBMCs from hospitalised psychiatric patients was first reported by Bode *et al.*, in 1994,⁹ and the detection of BDV RNA in PBMCs of neonatally infected rats¹⁰ stimulated investigation and detection in psychiatric patients.¹¹ The detection of antibodies in humans was first reported in 1985, in patients with depression and psychiatric disorders.¹² The s-antigen (p40 and p24 complex) initiates the major humoral response, and has been the main target for serological studies.¹³

The hypothesis that BDV is associated with human clinical illness is controversial. However, should even a small fraction of mental illness be attributable to infection with BDV, this would be an important finding, not least because illness in that sub-population would be potentially preventable. We performed a systematic review¹⁴ to: evaluate the evidence that BDV infects humans; evaluate the role of BDV in human neuropsychiatric illness; and assess the suitability of currently available laboratory methods for human epidemiological studies.

Methods

On-line and library databases were used to identify any studies involving the detection of BDV or its markers in human subjects. The following sources were searched using the term 'Borna': Medline, 1966 to January 2000; Institute for Science Information (Current Contents and Science Citation Index); British Library Document Supply Centre (including System for Information on Grey Literature); BIDS, 1981 to January 2000; BIOSYS via Datastar on-line index system, 1970 to January 2000; and the Cochrane Collaboration on CD-Rom.

The search was not limited by language or sources. Inclusion criteria were studies published prior to January 2000, where the study participants = human, investigation = antibody or antigen or viral markers, outcomes = prevalence data. This was followed by hand searching of the literature for further references, and by consultation with leading researchers, facilitated by the first UK workshop on Borna Disease Virus in March 2000.¹⁵

The primary studies were identified and assessed by the same researcher (RC). Due to the nature of the investigations and variation in methods and study groups found, meta-analysis was not undertaken. However, results were summarized to describe the studies undertaken, show the groups investigated, the tissue tested, laboratory methods used and results obtained. To facilitate cross-referencing, reports were allocated a study number, generated in chronological order. Seroprevalence (antibody) studies were allocated a number prefixed with 'Ab', studies for the prevalence of viral antigens or genetic markers were numbered with a 'V' prefix, and studies for both, were given an 'AbV' prefix. Individual case studies were described separately.

Results

We identified 75 publications that contained details of 50 human case or prevalence studies of Borna disease. The study designs identified were: five case studies of individual patients, and 44 (sero)prevalence studies of 138 groups of patients, including five studies of monozygotic twins (Tables 1–3). In 24/44 (55%) studies covering 64 groups of patients, a control or a comparison group was also tested to compare prevalence and to measure association between markers for infection and clinical illness. Epidemiological and exposure data was additionally gathered in just two studies. There were no studies in which the study design described the random or systematic recruitment of cases, selection of controls or systematic recruitment of a study cohort.

Case studies

A case of encephalitis with neurological and psychiatric symptomology was attributed to BDV when an IgG titre of 1/10 was detected.⁸⁴ Two family cases of psychiatric disorders were detected among 200 seropositive psychiatric cases. In both families, in addition to the primary patient, a second family member also had psychiatric or neurological illness as well as BDV antibodies.⁸⁵ BDV antibodies

Table 1 Published Borna disease virus (sero)prevalence studies

Study code	Study description	References
Ab 1	Prevalence of BDV antibodies in patients with unipolar and bipolar depression in Philadelphia (US) and patients with heterogenous psychiatric disorders in Giessen and Wurzburg (Germany), and in local control subjects.	12, 16
Ab 2	Prevalence of BDV antibodies in patients with HIV, EBV, affective disorder, drug abusers and a control group in US and West Germany.	17
Ab 3	Prevalence of BDV antibodies in psychiatric in-patients, and newly admitted psychiatric, neurological and surgery patients in Germany. MRI scans of psychiatric and surgical patients. CSF testing of further seropositive psychiatric patients, including virus isolation and infectivity in rabbits from two neurological and one psychiatric patient.	18–24
Ab 4	Retrospective prevalence of antibodies to BDV S-antigen amongst different groups of psychiatric, neurological and chronically infected people in different countries and in a control group from the population and blood donors in the US and Europe.	25, 26
Ab 5	Prevalence of antibodies to BDV S-antigen in patients with chronic fatigue syndrome in Germany.	27
Ab 6	Prevalence of antibodies to BDV 24 kDa and 38/40 kDa proteins in affective disorder patients and local healthy comparison subjects in Pennsylvania, USA.	28, 29
Ab 7	A longitudinal prevalence study of antibodies to BDV 38/40 kDa proteins in psychiatric patients in Germany.	30–32
Ab 8	Prevalence of BDV antibodies to gp18 and p40 in people exposed to farmed ostriches and a matched control group of blood donors in Israel.	33
Ab 9	Prevalence BDV antibodies in schizophrenic patients and community controls in Maryland, US and further investigation of deficit syndrome positive and negative patients.	34, 35
Ab 10	Prevalence of BDV antibodies in schizophrenic patients in Pennsylvania, US.	36
Ab 11	Prevalence of BDV antibodies in multiple sclerosis patients in Germany.	37
Ab 12	Prevalence of antibodies to BDV p24 in various HIV infected groups and AIDS patients in Thailand.	38
Ab 13	Prevalence of BDV antibodies in monozygotic twins in the US concordant and discordant for schizophrenia, bipolar depression and other psychoses.	39, 40
Ab 14	Prevalence of BDV antibodies in first episode schizophrenics in the US and seroconversion at 6-month follow-up, and comparison with unspecified controls.	41, 42
Ab 15	Prevalence of antibodies to BDV p40 in schizophrenic patients and healthy medical staff in Japan.	43
Ab 16	Prevalence of BDV antibodies in patients with panic disorder and healthy controls in Vienna, Austria.	44
Ab 17	Prevalence of antibodies to BDV p24 and p40 in Chinese schizophrenic patients, family members, mental health workers and non-psychiatric controls in Taiwan.	45
Ab 18	Prevalence of BDV antibodies in veterinarians practising in Styria, Austria.	46–48
Ab 19	Prevalence of BDV antibodies in patients with a variety of psychiatric, neurological and other disorders and blood donors in Japan.	49

were detected in a patient with motor neurone disease at a titre of 1/20.⁸⁶ Six blood samples were taken from a patient with somatization disorder and schizophrenia, and whole blood tested for p24 and p40 gene sequences by reverse transcriptive polymerase chain reaction (RT-PCR).⁸⁷ All samples contained both gene sequences.

Prevalence studies

Nineteen of 44 (43%) studies investigated seroprevalence, 11/44 (25%) investigated the prevalence of BDV, and 14/44 (32%) investigated both. Seroprevalence studies have been conducted in USA (5 studies), Germany (3 studies), Austria (2 studies),

Table 2 Summary of seroprevalence and prevalence of Borna disease virus in different study groups

Study group type	Seroprevalence		BDV prevalence	
	Positive studies/total studies	Median (range)	Positive studies/total studies	Median (range)
Unspecified psychiatric in-patients	9/11	7% (0–30%)	6/7	11% (0–54%)
Schizophrenia	10/13	5% (0–45%)	5/13	0% (0–82%)
Major depressive disorder	6/6	4% (1–12%)	1/8	0% (0–40%)
Neurotic, personality, adjustment and mood disorder	3/4	6% (0–15%)	1/1	4% (4%)
Mental disorders relating to alcohol and drugs	2/2	4% (2–5%)		NT
Mental retardation	0/2	0% (0%)		NT
Neurological patients	3/6	1% (0–5%)	1/2	1% (0–2%)
Dementia	0/3	0% (0%)	0/1	0% (0%)
Alzheimer's disease	1/1	1% (1%)	1/3	0% (0–11%)
Multiple sclerosis	1/3	5% (0–13%)	0/1	0% (0%)
Epilepsy	1/1	1% (1–1%)		NT
Parkinson's disease		NT	1/2	8% (0–17%)
Brain tumour		NT	1/1	14% (14%)
Chronic fatigue syndrome	2/4	5% (0–34%)	3/4	12% (0–24%)
HIV/AIDS	8/8	11% (1–48%)		NT
EBV	2/3	1% (0–6%)		NT
Parasitological infections	4/4	8% (5–19%)		NT
Occupational groups	2/3	10% (0–46%)		NT
Other groups	9/13	7% (0–36%)	3/8	0% (0–7%)

NT, none tested.

Table 3 Published studies of the prevalence of viral markers for Borna disease virus

Study code	Study description	Reference
V 1	Prevalence of p40 and p24 gene coding sequences in brain tumour patients in Japan.	50
V 2	Prevalence of p40 antigen and p40 gene coding sequences in brains of patients with histopathological characteristics on an Alzheimer Disease database in California, US.	51
V 3	Prevalence of P gene (p24) coding sequences in brain samples from neuropsychiatric patients and normal controls in North America and Europe.	52
V 4	Prevalence of P gene (p24) coding sequences in brain samples from neuropsychiatric patients and controls in Japan.	53
V 5	Prevalence of p24 gene coding sequences in normal brains at post mortem in Japan.	54
V 6	Prevalence of p40 gene coding sequences in whole peripheral blood and PBMCs from psychiatric patients in various regions of Germany with controls from one region.	55
V 7	Prevalence of p40 gene coding sequences in peripheral blood from blood donors and immunocompromised heart transfusion patients in Germany.	56
V 8	Prevalence of p24 gene coding sequences in PBMCs from patients with mood disorders or schizophrenia, and from age and sex matched blood donors, in Japan.	57
V 9	Prevalence of p40 gene coding sequences in brain, CSF and PBMCs of schizophrenics in US.	58
V 10	Prevalence of p24 gene coding sequences in PBMCs of psychiatric patients with schizophrenia or bipolar disorder in Korea.	59
V 11	Prevalence of unspecified virus gene coding sequences in brain tissue from neuropsychiatric and medical patients and healthy subjects at post mortem.	60

Japan (2 studies), Taiwan, Thailand and Israel (1 study each), and there were multicentre studies involving USA and Germany (2 studies), and USA, Europe and Africa (1 study). The location of one study was not specified. Prevalence studies have been conducted in Japan (4 studies), USA (3 studies), Germany (2 studies), Korea (1 study), and in USA and Europe as a multicentre study. Studies of both seroprevalence and prevalence have been conducted in Japan (8 studies), Germany (3 studies),

UK (1 study), the Netherlands (1 study), and in one multicentre study in USA and Germany. Studies were predominantly of adults, but children were included in one study.

Prevalence in patient groups

A variety of patient groups have been studied (Tables 4 and 5): unspecified psychiatric patients, schizophrenia cases, major depressive disorder,

Table 4 Published studies combining seroprevalence and prevalence of Borna disease viral markers

Study code	Study description	References
AbV 1	Detection of BDV antigens in populations of blood monocytes by flow cytometry in samples from psychiatric patients. Detection of p24 and p40 gene coding sequences in selected antigen positive blood samples, and virus isolated and characterized from further selected samples.	9, 11, 61–63
AbV 2	Prevalence of p24 gene coding sequence in blood donors in two areas of Japan (Sapporo and Tokyo), and prevalence of antibodies to BDV p24 in the Sapporo blood donors.	64
AbV 3	Prevalence of p24 gene coding sequence and prevalence of antibodies to BDV p24 in Japanese psychiatric patients. Sequence variability investigated in cDNA cloned and sequenced from three patients.	65, 66
AbV 4	Prevalence of antibodies to BDV p24 in Japanese CFS patients and healthy controls, and prevalence of p24 gene coding sequences in a further group of CFS patients. (Preliminary studies were for prevalence and seroprevalence in a sub-cohort of 25 CFS patients.)	67, 68
AbV 5	Prevalence of p24 gene coding sequence and prevalence of BDV antibodies in psychiatric patients and blood donors in Kumamoto, Japan.	69
AbV 6	Prevalence of antibodies to BDV 16 kDa, 24 kDa and 40 kDa antigens in psychiatric patients and controls in Hamburg, Germany, and prevalence of p24 and p40 gene coding sequences in a subset of those tested for antibodies.	70
AbV 7	Prevalence of antibodies to BDV p38 in psychiatric patients and controls in Japan and prevalence of p24 gene coding sequences in a subset of these.	71
AbV 8	Prevalence of BDV antibodies and prevalence of p24 gene coding sequence in schizophrenic patients and normal controls in Philadelphia.	72
AbV 9	Prevalence of BDV antibodies and prevalence of p24 gene coding sequences in British CFS patients, those with affective disorders and normal controls. Brain tissue examined for virus.	73, 74
AbV 10	Prevalence of antibodies to BDV p24 and p40 and prevalence of p24 coding sequences in Japanese schizophrenic patients and healthy controls. Some epidemiological data also collected and analysed.	75–78
AbV 11	Prevalence of BDV antibodies and prevalence of BDVp24 and p40 gene coding sequences in CSF of depressive patients and MS patients and healthy controls in Munich, Germany.	79, 80
AbV 12	Prevalence of antibodies to BDV and prevalence of BDV p40 gene coding sequences in Alzheimer and vascular dementia patients and healthy blood donor controls.	81
AbV 13	Only prevalence of BDV antibodies results given.	82
AbV 14	Prevalence of antibodies to BDV p24 in blood donors in towns near Thoroughbred horse farms. 7 antibody positive donors also tested for BDV RNA. Prevalence of BDV p24 gene coding sequence also tested in one group of blood donors.	83

Table 5 Prevalence of antibodies to Borna disease virus by study group

Location	Study number and study group description	Study group seroprevalence	Test sample	Test method	Control group sero-prevalence	Quoted <i>p</i> value
	<i>Unspecified psychiatric in-patients</i>					
Japan	AbV 3	18/60 (30%)	Blood plasma	Western blot		
	Unspecified psychiatric in-patients					
Japan	AbV 5	13/33 (24%)	Blood plasma	Indirect IFA		
	Unspecified psychiatric in-patients					
Germany	AbV 6	7/34 (21%)	Blood sera	Western blot	3/203 (2%)	
	Mental disorders (organic factors)					
Germany	AbV 1	12/68 (18%)	Blood sera	Double stain IFA		
	Unspecified psychiatric in-patients					
Germany	AbV 6	2/20 (10%)	Blood sera	Western blot	3/203 (2%)	
	Unspecified mental disorders on admission					
Germany	Ab 3	68/1003 (7%)	Blood sera	Indirect IFA		
	Unspecified psychiatric in-patients					
Germany	Ab 3	140/2377 (6%)	Blood sera	Indirect IFA		
	Newly admitted psychiatric in-patients					
Germany	AbV 11	2/128 (2%)				
	Unspecified psychiatric in-patients					
Germany	Ab 1	4/694 (1%)	Blood sera	Indirect IFA	0/95 (0%)	
	Unspecified psychiatric in-patients					
Japan	Ab 19	0/164 (0%)	Blood sera	ECL immunoassay		
	Unspecified psychiatric in-patients					
Japan	AbV 7	0/44 (0%)	Blood plasma	Indirect IFA	0/70 (0%)	
	Unspecified psychiatric in-patients					
	<i>Schizophrenic patients</i>					
Japan	AbV 10	30/67 (45%)	Blood plasma	Western blot	0/26 (0%)	
	Schizophrenics					
USA	Ab 9	13/90 (14%)	Blood sera	Western blot	0/20 (0%)	0.06
	Schizophrenic outpatients					
Germany	AbV 6	16/114 (14%)	Blood sera	Western blot	3/203 (2%)	
	Schizophrenia and other psychotic disorders					

USA	Ab 14	10/82 (12%)	Blood sera	Western blot	0/22 (0%)	
	First episode schizophrenia					
Taiwan	Ab 17	38/314 (12%)	Blood sera	Western blot	8/277 (3%)	
	Chronic schizophrenia					
Netherlands	AbV13	3/27 (11%)	Not stated	IFA	6/29 (21%)	
	Male schizophrenic immigrants					
USA	Ab 10	2/41 (5%)	Blood sera	Western blot		
	Schizophrenics					
USA	Ab 13	1/26 (4%)	Blood sera	Western blot		
	Monozygotic twins					
	concordant schizophrenia					
Japan	Ab 19	26/845 (3%)	Blood sera	ECL immunoassay	10/917 (1%)	<0.005
	Schizophrenics					
Japan	Ab 15	2/70 (3%)	Blood sera	ELISA	1/40 (3%)	
	Chronic schizophrenia					
USA	Ab 2	0/49 (0%)	Blood sera	Double stain IFA	11/540 (2%)	
	Schizophrenics					
Japan	AbV 7	0/179 (0%)	Blood plasma	Indirect IFA	0/70 (0%)	
	Schizophrenics					
USA	AbV 8	2/10 (2%)	Blood sera	Western blot	0/10 (0%)	
	Schizophrenics					
	<i>Major depressive disorder</i>					
Germany	AbV 6	6/52 (12%)	Blood sera	Western blot	3/203 (2%)	
	Affective psychoses and disorder					
USA	Ab 6	9/138 (7%)	Blood sera	Western blot	1/117 (1%)	<0.025 q
	Major depressive disorder					
USA	Ab 1	12/285 (4%)	Blood sera	Indirect IFA	0/105 (0%)	
	Unipolar and bipolar depression					
USA	Ab 4	7/187 (4%)	Blood sera	Double stain IFA	11/483 (2%)	
	Bipolar depression					
USA	Ab 2	13/593 (2%)	Blood sera	Double stain IFA	11/540 (2%)	
	Depression					
USA	Ab 4	5/363 (1%)	Blood sera	Double stain IFA	11/483 (2%)	
	Unipolar depression					
	Neurotic, personality, adjustment					
	and mood disorder					
Germany	AbV 6	8/54 (15%)	Blood sera	Western blot	3/203 (2%)	
	Neurotic, personality,					
	adjustment disorder					

(Continued)

Table 5 Continued

Location	Study number and study group description	Study group seroprevalence	Test sample	Test method	Control group sero-prevalence	Quoted <i>p</i> value
Austria	Ab 16 Outpatients with panic disorder	4/55 (7%)	Not stated	Not stated	2/34 (6%)	>0.05 <i>q</i>
Japan	Ab 19 Mood disorder	9/251 (4%)	Blood sera	ECL immunoassay	10/917 (1%)	<0.05 <i>q</i>
Japan	AbV 7 Mood disorder	0/123 (0%)	Blood plasma	Indirect IFA	0/70 (0%)	
<i>Mental disorders relating to alcohol and drugs</i>						
Germany	AbV 6 Alcohol/drug related mental disorders	1/22 (5%)	Blood sera	Western blot	3/203 (2%)	
Japan	Ab 19 Alcohol addiction	1/42 (2%)	Blood sera	ECL immunoassay		
<i>Mental retardation</i>						
Japan	Ab 19 Mental retardation	0/25 (0%)	Blood sera	ECL immunoassay		
Germany	AbV 6 Mental retardation	0/2 (0%)	Blood sera	Western blot		
<i>Neurological patients</i>						
Germany	Ab 3 Newly admitted unspecified patients	87/1791 (5%)	CSF	Indirect IFA		
USA	Ab 4 Peripheral neurology	4/92 (4%)	Blood sera	Double stain IFA	11/483 (2%)	>0.05 <i>q</i>
Germany	AbV 11 Neurological inpatients	1/102 (1%)	CSF	Indirect IFA	None	
Germany	Ab 4 Purulent non-viral encephalitis	0/54 (0%)	Blood sera	Double stain IFA		
Japan	Ab 19 Encephalitis	0/33 (0%)	Blood sera	ECL immunoassay		
Japan	Ab 19 Degenerative diseases	0/14 (0%)	Blood sera	ECL immunoassay		

<i>Dementia</i>						
Japan	AbV 12	0/10 (0%)	Blood plasma	Indirect IFA	4/36 (11%)	
Japan	Vascular dementia patients Ab 19	0/46 (0%)	Blood sera	ECL immunoassay		
Japan	Vascular dementia AbV 12	0/10 (0%)	Blood plasma	Indirect IFA	4/36 (11%)	
	Dementia patients					
<i>Alzheimer's disease</i>						
Japan	Ab 19	1/89 (1%)	Blood sera	ECL immunoassay		
	Alzheimer's disease					
<i>Multiple sclerosis</i>						
Germany	Ab 4	15/114 (13%)	Blood sera	Double stain IFA	11/483 (2%)	<0.0001 <i>q</i>
	Multiple sclerosis					
Germany	Ab 11	0/50 (0%)	CSF and blood sera	Indirect IFA		
	Multiple sclerosis					
Japan	Ab 19	0/67 (0%)	Blood sera	ECL immunoassay		
	Multiple sclerosis					
<i>Epilepsy</i>						
Japan	Ab 19	3/214 (1%)	Blood sera	ECL immunoassay		
	Epilepsy					
<i>Chronic fatigue syndrome</i>						
Japan	AbV 4	30/89 (34%)	Blood plasma or sera	ELISA	1/100 (1%)	<0.001
	Chronic fatigue syndrome					
UK	AbV 9	2/21 (10%)	Blood sera	Western blot	0/13 (0%)	
	Chronic fatigue syndrome					
Germany	Ab 5	0/50 (0%)	Blood sera	Double stain IFA	None	
	Chronic fatigue syndrome					
Japan	Ab 19	0/75 (0%)	Blood sera	ECL immunoassay		
	Chronic fatigue syndrome					
<i>HIV/AIDS</i>						
Thailand	Ab 12	24/50 (48%)	Blood sera	ELISA	2/103 (2%)	
	HIV Clade E with unspecified STD					
Thailand	Ab 12	12/67 (18%)	Blood sera	ELISA	None	
	AIDS patients					
Thailand	Ab 12	9/60 (15%)	Blood sera	ELISA	None	
	Asymptomatic HIV carriers					
Germany	Ab 4	34/244 (14%)	Blood sera	Double stain IFA	11/483 (3%)	
	HIV lymphadenopathy					

(Continued)

Table 5 Continued

Location	Study number and study group description	Study group seroprevalence	Test sample	Test method	Control group sero-prevalence	Quoted p value
Germany	Ab 4 HIV antibody peak	61/751 (8%)	Blood sera	Double stain IFA	11/483 (3%)	
Germany	Ab 2 HIV antibody	36/460 (8%)	Blood sera	Double stain IFA	11/540 (2%)	
Germany	Ab 4 HIV antibody start	12/273 (4%)	Blood sera	Double stain IFA	11/483 (3%)	
Japan	Ab 19 HIV including AIDS	1/85 (1%)	Blood sera	ECL immunoassay		
<i>Epstein Barr syndrome</i>						
Germany	Ab 4 Children	9/160 (6%)	Blood sera	Double stain IFA	11/483 (3%)	
Germany	Ab 4 Adults	1/89 (1%)	Blood sera	Double stain IFA	11/483 (3%)	
Germany	Ab 2 EBV	0/24 (0%)	Blood sera	Double stain IFA		
<i>Parasitological infections</i>						
East Africa	Ab 4 Children with schistosomiasis and malaria	9/48 (19%)	Blood sera	Double stain IFA	11/483 (3%)	<0.0001
East Africa	Ab 4 Children with schistosomiasis or malaria	1/12 (8%)	Blood sera	Double stain IFA	11/483 (3%)	<0.05
East Africa	Ab 4 Adults with schistosomiasis and malaria	10/145 (7%)	Blood sera	Double stain IFA	11/483 (3%)	<0.01
East Africa	Ab 4 Adults with schistosomiasis or malaria	4/81 (5%)	Blood sera	Double stain IFA	11/483 (3%)	<0.05
<i>Occupational groups</i>						
Israel	Ab 8 Workers exposed to ostriches	19/41 (46%)	Blood sera	ELISA	4/41 (10%)	<0.0001
Taiwan	Ab 17 Mental health workers	8/82 (10%)	Blood sera	Western blot		
Austria	Ab 18 Veterinarians	0/137 (0%)	Blood sera	Indirect IFA		

Other groups					
USA	Ab 13 Twenty five pairs of monozygotic twins discordant for schizophrenia	9/25 (36%) affected subjects 5/25 (20%) unaffected subjects	Blood sera	Western blot	
Taiwan	Ab 17 Family members of schizophrenic patients	16/132 (21%)	Blood sera	Western blot	
Germany	Ab 7 Sequentially tested psychiatric in-patients	14/71 (20%)	Blood sera	Double stain IFA	8/22 (36%) major depression
Japan	AbV 14 Blood donors from horse rearing area	36/428 (8%)	Blood sera	ELISA	1/100 (1%) <0.05
Japan	Ab 19 Multi-transfused patients	3/66 (5%)	Blood sera	ECL immunoassay	
Germany	Ab 2 HIV-antibody-negative drug abusers	4/106 (4%)	Blood sera	Double stain IFA	2/125 (2%) HIV negative non-drug abusers
Germany	Ab 3 Newly admitted surgery patients	20/569 (4%)	Blood sera	Indirect IFA	
Japan	Ab 19 Ocular diseases	19/1393 (1%)	Blood sera	ECL immunoassay	10/917 (1%) >0.05
Japan	AbV 2 Blood donors	1/100 (1%)	Blood sera	ELISA	
USA	Ab 13 Eight pairs monozygotic twins discordant for bipolar disorder	0/8 (0%) affected subjects 0/8 (0%) unaffected subjects	Blood sera	Western blot	
USA	Ab 13 Eight pairs normal monozygotic twins	0/16 (0%)	Blood sera	Western blot	
Japan	Ab 19 Autoimmune disease	0/50 (0%)	Blood sera	ECL immunoassay	
Japan	Ab 19 Leprosy	0/17 (0%)	Blood sera	ECL immunoassay	

other psychiatric disorders, neurology patients, dementia/Alzheimer's disease, multiple sclerosis, epilepsy and chronic fatigue syndrome, HIV/AIDS, EBV, parasitological infections and occupational groups. Five groups of monozygotic twins, concordant and discordant for schizophrenia and other disorders, have been studied. Patient groups were defined by internationally recognized criteria (Research Diagnostic Criteria, ICD codes, DSM codes or CDC guidelines) in 18/44 (41%) studies. Control or comparison groups were either from the local community or blood donors.

Prevalence and seroprevalence varied widely (Table 6) from 0% (no evidence of infection), while the highest seroprevalence in a single study group was 48% in 50 HIV Clade E patients with STDs in Thailand. The highest prevalence of virus was 82% in 11 brains of schizophrenic patients. However this is an outlying result, and the median for this patient group was 0%. Median seroprevalence figures were highest (11%) in HIV/AIDS patients, and the highest median prevalence was 13% in astrocytic brain tumours, although this was a single study.

Laboratory methods

Seroprevalence was mainly investigated by the detection of antibodies in serum or plasma (28 studies) but also in CSF and sera (2 studies), CSF only (1 study) and unspecified material (2 studies) (Table 4). Studies for genetic markers and viral antigens have been sought mainly in peripheral blood and PBMCs or leukocytes (PBLs) (15 studies), brain tissue (7 studies), CSF (1 study) and CSF, brain and blood (2 studies) (Table 5).

The laboratory methods used to test for BDV antibodies were: indirect immunofluorescence antibody/assay (IFA) (10 studies), double stain IFA (5 studies), Western blot using viral proteins (6 studies), Western blot using recombinant proteins (4 studies), enzyme-linked immunosorbent assay using recombinant proteins (recombinant ELISA) (4 studies), recombinant reverse type (RT) sandwich ELISA (1 study), electrochemiluminescence assay (1 study), and direct IFA (1 study); in one study, the method was not stated.

Test methods for antigens or genetic markers were immunohistochemistry with nested PCR (1 study), flow cytometry (1 study), enzyme immunoassay (EIA) (1 study), EIA with nested RT-PCR (1 study), and RT-PCR (3 studies). The most frequently used was nested RT-PCR for p24 and/or p40 gene sequences (17 studies).

Issues of sensitivity and specificity, reproducibility and reliability were rarely addressed.

A second test was usually used for confirmation of antibody tests (Western blotting), and genetic markers were confirmed by Southern blotting, immunohistochemistry, competition experiments, *in-situ* hybridization and direct sequencing.

Discussion

The neurological or psychiatric symptoms of human patients studied for Borna disease are clinically diverse, and given the frequently multifactorial aetiology of mental illness, an alternative explanation would be possible for all of these cases. The presence of markers of BDV infection in a clinically ill patient cannot infer causation (see reference 88). Indeed, individuals may be predisposed to BDV infection because of altered immunocompetence during repeated affective episodes.¹² Clinical case studies, however, are valuable for generating hypotheses that should be tested further in systematic epidemiological studies. These patients also provide the opportunity for molecular studies and characterisation of virus isolates.

Seroprevalence studies provide information and evidence of exposure to BDV, while prevalence studies measure the proportion of current infection. Many such studies have been carried out, in a variety of patient groups most usually selected by a clinical illness. However, while simple in their design, the validity of these studies is dependent on the sensitivity, specificity and reproducibility both of the diagnostic tests performed and of the diagnostic criteria used in the classification of patients. A wide range of seroprevalence even within specific clinical groups has been demonstrated, which may reflect geographical variation in prevalence, selection bias, or the different laboratory methods used. Median seroprevalence was highest for unspecified psychiatric in-patients (7%), neurotic, personality adjustment and mood disorder patients (6%), schizophrenics (5%) and patients with chronic fatigue syndrome (5%). These data might indicate a direction for future epidemiological studies. It is also of note that at least one person tested seropositive in every clinical group examined, except for those patients suffering from mental retardation and dementia. Other groups have been examined, probably opportunistically, such as children and adults with parasitological infections and HIV patients. Any other rationale for screening such patient groups is unclear.

Prevalence studies are usually of limited value in aetiological research, since prevalence reflects both the incidence and probability of surviving the disease.⁸⁸ In these studies, incident mental illness

is the outcome and previous exposure to BDV is the exposure of interest. To conclude that BDV has a role in any of these clinical syndromes requires that the prevalence in these groups (cases) be compared with the prevalence in another 'healthy' or 'non-case' group (controls). The outcome of such a study would be a measure of the strength of association between having a marker for BDV exposure and/or infection and being in a certain clinical group, conventionally expressed as an odds ratio. However, the odds ratio calculated in a case-control study will be a biased estimate unless the controls selected are representative of the population that are well, but could become cases. We have carried out a quantitative systematic review but have not attempted any meta-analysis methods (see reference 89). As it was unlikely that the same disease diagnostic criteria were applied across studies, and because of the variation in selection of controls, no attempt has been made to combine data from individual studies to calculate summary odds ratios. In many published BDV seroprevalence studies, there was not enough information provided on the selection of controls for the reader to assess whether they represented well individuals belonging to at-risk populations. In some studies the 'control' group should be considered a 'comparison' group, and was often small. In most studies, odds ratios were not calculated. Chen *et al.*⁴⁵ reported a positive association between presence of BDV antibodies and schizophrenia in Chinese patients from Taiwan and presented their results as an odds ratio (odds ratio 4.63, 95%CI 2.25–9.52).

Few studies have attempted to investigate possible sources of human exposure to BDV. One published study³³ found high seroprevalence in those exposed to ostriches (46%), but did not investigate possible confounding factors that may be associated both with handling ostriches and exposure to BDV, such as contact with other farm animals. Takahashi *et al.*⁸³ found a high seropositive rate in people living near horse farms. For some individual cases, animal exposure histories have been reported together with the results of the sampling of animal contacts (e.g. reference 84). Histories of exposure to seropositive animals or animals with a history of Borna disease is important in assessing putative human cases. Further epidemiological studies could be carried out to identify possible exposures to BDV. Further animal studies would also prove useful to establish the host range and assess possible modes of transmission of the virus, including vectors or reservoirs of infection. Also of interest would be any links between reported human cases. Chen *et al.*⁴⁵ also found higher BDV seroprevalence in

first degree relatives of schizophrenic patients and mental health workers. This raises the possibility of person-to-person transmission, but could also be the result of bias in selection of controls and/or confounding factors. Family clusters of Borna disease have been reported from Germany⁸⁴ and the identification and follow-up of such clusters could prove useful in providing further information on the zoonotic potential of BDV. Cases in the published literature attributed to infection with BDV have been from the geographical areas in central Europe where Borna disease is endemic in animals. Whilst this would appear consistent with the hypothesis that Borna disease is a zoonosis, it may simply be the result of ascertainment or publication bias.

There does appear to be evidence that people are exposed to BDV, or at least have antibodies that are reactive to BDV antigens. However, while a number of seroprevalence studies have been done, it is still difficult to conclude much about which population groups are most at risk and the natural history of human infection. Longitudinal studies are necessary to follow the serological response and clinical course in patients identified as antibody positive. Evidence that humans are infected with BDV, or a BDV-like virus, rather than exposed, is even more controversial. Conflicting results have arisen and evidence of infection or exposure by the detection of viral markers or antibodies, is not consistently linked to one currently defined disease in humans. Some results show that BDV RNA is detected in the absence of antibodies in serum, further complicating matters.

An agreed gold standard diagnostic test for BDV is lacking, and there is speculation concerning the specificity of the tests currently used. It has been suggested that PCR positives are the result of laboratory contamination.^{90,91} Further investigations have revealed novel isolates that may not amplify with previously described markers, and variation may be greater than previously thought.⁹² Multi-centre studies have tested the same samples in a number of different laboratories by a number of different methods but this has been performed in order to confirm a result rather than to validate tests in terms of their sensitivity and specificity, reproducibility and reliability. Of course, it is not possible to validate tests using populations of known cases and controls, because is not yet clear what criteria should be used to define a human case. A well-validated blood test would appear to be the most suitable test for epidemiological studies. Certainly there are practical difficulties in using brain tissue or CSF. In the absence of consensus as to the validity, it would appear that an antibody test

Table 6 Prevalence of Borna disease virus, by study group

Location	Study number, study group description	Tissue tested	Prevalence in study group	Test method	Prevalence in control group
<i>Unspecified psychiatric in-patients</i>					
Germany	AbV 1 Chronic mental illness	PBMCs	37/68 (54%)	Flow cytometry	
Germany	AbV 6 A subset of psychiatric patients	PBMCs	13/26 (50%)	Nested RT-PCR	0/23 (0%)
Japan	AbV 3 Psychiatric inpatients	PBMCs	22/60 (37%)	Nested RT-PCR	
Japan	AbV 5 Psychiatric patients	PBMCs	6/55 (11%)	Nested RT-PCR	
Germany	AbV 11 Psychiatric inpatients	CSF	3/128 (2%)	EIA	
Japan	AbV 7 A subset of psychiatric patients	PBMCs	2/106 (2%)	RT-PCR	0/12 (0%)
Germany	V 6 Psychiatric disorders	Whole blood	0/39 (0%)	Nested RT-PCR	
<i>Schizophrenia and other psychotic disorders</i>					
Europe and US	V 3 Schizophrenic	Brain	9/11 (82%)	Nested RT-PCR	0/10 (0%)
Japan	V 4 Schizophrenic	Brain	3/9 (33%)	Nested RT-PCR	2/31 (7%)
Japan	AbV 10 Schizophrenia patients	Whole blood	6/67 (9%)	Nested RT-PCR	
Japan	V 8 Schizophrenic	PBMCs	3/77 (4%)	Nested RT-PCR	
Germany	V 6 Schizophrenic	Whole blood	0/59 (0%)	Nested RT-PCR	
Germany	V 6 Schizophrenic	Whole blood	0/10 (0%)	Nested RT-PCR	
Germany	V 6 Schizoaffective disorder	Blood/PBMCs	0/20 (0%)	Nested RT-PCR	
US	V 9 Schizophrenic	CSF	0/48 (0%)	Cell culture and nested RT-PCR EIA	

US	V 9	Brain	0/3	Nested RT-PCR EIA	
Europe and US	Schizophrenic V 3	Brain	0/2	Nested RT-PCR	
Korea	Psychotic disorder V 10	Not stated	0/39 (0%)	Nested RT-PCR	
US	Schizophrenic V 11	Brain	0/16 (0%)	RT-PCR	
US	Schizophrenic AbV 8 Schizophrenic patients	PBLs	2/10 (20%) in 1 lab, 0/10 (0%) in 2 other labs	RT-PCR	
Major depressive disorder Europe and US	V 3	Brain	2/5 (40%)	Nested RT-PCR	0/10 (0%)
Europe and US	Bipolar disorder V 3	Brain	0/6 (0%)	Nested RT-PCR	0/10 (0%)
Germany	Major depressive disorder V 6	Whole blood/PBMCs	0/41 (0%)	Nested RT-PCR	
Germany	Major depressive disorder V 6	Whole blood/PBMCs	0/10 (0%)	Nested RT-PCR	
US	Bipolar disorder V 9	CSF/Brain/PBMCs	0/33 (0%)	Nested RT-PCR	
Korea	Bipolar disorder V 10	Not stated	0/9 (0%)	Nested RT-PCR	
US	Major depression V 11	Brain	0/5 (0%)	RT-PCR	
UK	Affective disorder AbV 9 Severe depression	PBMCs	0/8 (0%)	Nested RT-PCR	
Neurotic, personality, adjustment and mood disorder Japan	V 8	PBMCs	2/49 (4%)	Nested RT-PCR	
Germany	Mood disorder patients Neurological patients AbV11	CSF	2/102 (2%)	EIA	
US	Neurological inpatients V 11 Neurological disorder	Brain	0/17 (0%)	RT-PCR	

(Continued)

Table 6 Continued

Location	Study number, study group description	Tissue tested	Prevalence in study group	Test method	Prevalence in control group
<i>Dementia</i> Japan	AbV 12 Vascular dementia patients Multiple sclerosis	PBMCs	0/10 (0%)	Nested RT-PCR	0/36 (0%)
Europe and US	V 3 Multiple sclerosis	Brain	0/5 (0%)	Nested RT-PCR	0/10 (0%)
<i>Alzheimer's disease</i> US	V 2 Alzheimer's disease	Brain	4/34 (12%)	Immunohistochemistry	0/2 (0%)
Europe and US	V 3 Alzheimer's disease	Brain	0/19 (0%)	Nested RT-PCR	0/10 (0%)
Japan	AbV 12 Alzheimer's disease	PBMCs	0/10 (0%)	Nested RT-PCR	0/36 (0%)
<i>Parkinson's disease</i> Japan	V 4 Parkinson's disease	Brain	1/6 (17%)	Nested RT-PCR	2/31 (7%)
Europe and US	V 3 Parkinson's disease	Brain	0/11 (0%)	Nested RT-PCR	0/10 (0%)
<i>Brain tumours</i> Japan	V1 Astrocytic tumours	Brain tumours	5/37 (14%)	Nested RT-PCR	None
<i>Chronic fatigue syndrome</i> Japan	AbV 4 Sub-cohort of CFS patients	PBMC	6/25 (24%)	Nested RT-PCR	

Japan	AbV4	PBMC	7/57 (12%)	Nested RT-PCR	
	Further cohort of CFS patients				
UK	AbV 9	CSF	1/10 (10%)	Nested RT-PCR	0/10 (0%) CSF
UK	Subset of CSF from CFS patients				
	AbV 9	Sera/PBMC	0/21 (0%)	Nested RT-PCR	0/31 (0%)
	CFS patients				
<i>Other groups</i>					
Japan	V 5	Brain	2/30 (7%)	Nested RT-PCR	
	Normal brains at post mortem				
Japan	AbV 2	PBMCs	5/100 (5%)	Nested RT-PCR	
	Blood donors				
US	AbV 2	PBMCs	3/72 (4%)	Nested RT-PCR	
	Blood donors				
Germany	V 7	Blood	0/200 (0%)	RT-PCR	
	Blood donors				
US	V 9	CSF	0/18 (0%)	Cell culture and nested RT-PCR EIA	
	Nine sets monozygotic twins discordant for Schizophrenia	PBMCs	0/18 (0%)		
US	V 11	Brain	0/27 (0%)	RT-PCR	
	Medical patients				
US	V 11	Brain	0/20 (0%)	RT-PCR	
	Healthy subjects				
Germany	V 7	Blood	0/500 (0%)	Nested RT-PCR	
	Immunosuppressed heart transplant patients				

would be the most suitable test by which to screen large population samples. A 'gold-standard' serological test for Borna disease virus should be developed that is suitable for testing animal and human samples in epidemiological studies. Clinical specimens should be swapped between experts, and data should be gathered on the sensitivity, specificity and reproducibility of any such 'gold-standard' developed. This might be most feasible using sera from clinical cases of Borna disease in animals.

The hypothesis that Borna disease is a zoonosis merits serious consideration and is worthy of further investigation.

Acknowledgements

This research was funded by the Health and Safety Executive (Project number: 3490/R52.118).

References

- Zwick W. Borna'sche Krankheit und enzephalomyelitis der Tiere. In: Gildemeister F, Hagen E, Waldmann O, eds. *Handbuch der Viruserkrankungen*, Vol. 2. Jena, VEB Fischer Verlag, 1939:254–354.
- Richt J, Pfeuffer I, Christ M, Frese K, Bechter K, Herzog S. Borna Disease Virus infection in animals and humans. *Emerg Infect Dis* 1997; **3**:343–52.
- Gosztonyi G, Ludwig H. Borna disease of horses. *Acta Neuropathol* 1984; **64**:213–21.
- Gonzalez-Dunia D, Sauder C, de la Torre JC. Borna Disease Virus and the brain. *Brain Res Bull* 1997; **44**:647–64.
- Hatalski CG, Lewis AJ, Lipkin WI. Borna Disease. *Emerg Infect Dis* 1997; **3**:129–35.
- Richt JA, Herzog S, Haberkettl K, Rott R. Demonstration of Borna disease virus-specific RNA in secretions of naturally infected horses by the polymerase chain reaction. *Med Microbiol Immunol* 1993; **182**:293–304.
- Herzog S, Frese K, Richt JA, Rott R. Ein Beitrag zur epizootiologie der Borna'schen Krankheit beim Pferd. *Wien Tierarztl Mschr* 1994; **81**:374–9.
- Cubitt B, de la Torre C. Borna Disease virus (BDV) a nonsegmented RNA virus, replicates in the nuclei of infected cells where infectious BDV ribonucleoproteins are present. *J Virol* 1994; **68**:1371–81.
- Bode L, Steinbach F, Ludwig H. A novel marker for Borna disease virus infection. *Lancet* 1994; **343**:297–8.
- Sierra-Honigmann AM, Rubin SA, Estafanous MG, Yolken RH, Carbone KM. Borna disease virus in peripheral blood mononuclear and bone marrow cells of neonatally and chronically infected rats. *J Neuroimmunol* 1993; **45**:31–6.
- Bode L, Zimmermann W, Ferszt R, Steinbach F, Ludwig H. Borna disease virus genome transcribed and expressed in psychiatric patients. *Nat Med* 1995; **1**:232–6.
- Amsterdam JD, Winokur H, Dyson W, Herzog S, Rott R, Koprowski H. Borna disease virus: a possible etiologic factor in human affective disorders? *Arch Gen Psychiatry* 1985; **42**:1093–6.
- Ludwig H, Furuya K, Bode L, Durrwald R, Lee DS. Biology and neurobiology of Borna disease virus (BDV), defined by antibodies, neutralizability and their pathogenic potential. *Arch Virol* 1993; **7**:111–33.
- Mulrow CD. Rationale for systematic reviews. *Br Med J* 1994; **309**:597–9.
- Anon. 1st UK Workshop on Borna Disease Virus, Rhondda, Wales, UK, March 2000. [<http://www.phls.wales.nhs.uk/bconproc.htm>].
- Rott R, Herzog S, Fleischer B, Winokur H, Amsterdam J, Dyson W, et al. Detection of serum antibodies to Borna disease virus in patients with psychiatric disorders. *Science* 1985; **228**:755–6.
- Bode L, Riegel S, Ludwig H, Amsterdam JD, Lange W, Koprowski H. Borna disease virus-specific antibodies in patients with HIV infection and with mental disorders. *Lancet* 1988; **ii**:689.
- Bechter K, Herzog S, Fleischer B, Schuttler R, Rott R. Magnetic resonance imaging in psychiatric patients, with and without serum antibodies against Borna Disease. *Der Nervenarzt* 1987; **58**:617–24.
- Bechter K, Herzog S, Schuttler R, Rott R. Die Borna'sche Krankheit-wahrscheinlich auch eine menschliche Krankheit:neue Ergebnisse. In: Saletu B, ed. *Biologische Psychiatrie*. Stuttgart, Thieme, 1989:17–21.
- Rott R, Herzog S, Bechter K, Frese K. Borna Disease, a possible hazard for man? *Arch Virol* 1991; **118**:143–9.
- Bechter K, Herzog S, Schuttler R. Possible significance of Borna Disease for humans. *Neurol Psychiat Brain Res* 1992; **1**:23–9.
- Bechter K, Bauer M, Hestler HC, Rott R. English title: MRI findings in Borna Disease Virus seropositive psychiatric patients and controls: and extended study. *Der Nervenarzt* 1994; **65**:169–74.
- Bechter K, Herzog S, Behr W, Schuttler R. Investigations of cerebrospinal fluid in Borna disease virus seropositive psychiatric patients. *Eurp Psychiat* 1995; **10**:250–8.
- Bechter K, Herzog S, Schuttler R. Borna Disease Virus (BDV)—possible cause of a spectrum of human neuropsychiatric disorders. *Pharmacopsychiat* 1997; **30**:151 (proc 20th symposium AGNP, Nuremberg 1997).
- Bode L, Riegel S, Czech G, Lange W, Ludwig H. Increased incidence of Borna Disease Virus infection in chronically diseased patients. Abstract, VIIIth Int Congress of Virol Berlin 26–29th August 1990.
- Bode L, Riegel S, Lange W, Ludwig H. Human infections with Borna Disease Virus: seroprevalence in patients with chronic diseases and healthy individuals. *J Med Virol* 1992; **36**:309–15.
- Bode L, Komaroff AL, Ludwig H. No serologic evidence of Borna Disease Virus in patients with Chronic Fatigue Syndrome. *Clin Infect Dis* 1992; **15**:1049.
- Amsterdam JD, Fu ZF, Dietzschold B. Anti-Borna Disease (BD) Virus-specific antibodies in depressed patients measured by Western immunoblot technique. *Biolog Psychiat* 1992; **31**:102A.
- Fu ZF, Amsterdam JD, Kao M, Shankar V, Koprowski H, Dietzschold B. Detection of Borna Disease virus-reactive antibodies from patients with affective disorders by Western immunoblot technique. *J Affect Dis* 1993; **27**:61–8.
- Bode L, Querfurth H, Ferszt R, Gosztonyi G, Rigas N, Czech G, et al. Antibodies to Borna disease virus are

- four times as frequent in depressives as in controls. In: *Wissenschaftswoche 1991, Forschungsprojekte am Klinikum Steglitz, Freie Universität Berlin*. Berlin, Freie Universität Berlin, 1991:269–71.
31. Bode L, Czech G, Ferszt R, Ludwig H. Bornavirus-Infektion beim Menschen: eine neue Zoonose? In: *Bericht des 4. Hohenheimer Seminars'Aktuelle Zoonosen, Sept 16–17, 1992, Stuttgart*. Giessen, Deutsche Veterinarmedizinische Gesellschaft (DVG), 1992:138–47.
 32. Bode L, Ferszt R, Czech G. Borna Disease virus infection and affective disorders in man. *Arch Virol* 1993; **7(suppl.)**: 159–67.
 33. Weisman Y, Huminer D, Malkinson M, Meir R, Kliche S, Lipkin WI, *et al.* Borna disease virus antibodies among workers exposed to infected ostriches. *Lancet* 1994; **344**: 1232–3.
 34. Waltrip RW, Buchanan RW, Summerfelt A, Breier A, Carpenter WT, Bryant NL, *et al.* Borna disease virus and schizophrenia. *Psych Res* 1995; **56**:33–44.
 35. Waltrip RW, Buchanan RW, Carpenter WT, Kirkpatrick B, Summerfelt A, Breier A, *et al.* Borna Disease Virus antibodies and the deficit syndrome of schizophrenia. *Schizophr Res* 1997; **23**:253–7.
 36. Pelle M, Alexander RC, Fu ZF, Dietzschold B, Koprowski H. Borna Disease Antibodies in Schizophrenia. *Schizophr Res* 1995; **15**:189.
 37. Kitze B, Herzog S, Rieckmann P, Poser S, Richt J. No evidence of Borna disease virus-specific antibodies in multiple sclerosis patients in Germany. *J Neurol* 1996; **243**:660–2.
 38. Auwanit W, Ayuthaya PIN, Nakaya T, Fujiwara S, Kurata T, Yamanishi K, *et al.* Unusually high seroprevalence of Borna Disease Virus in Clade E human immunodeficiency virus type 1-infected patients with sexually transmitted diseases in Thailand. *Clin Diag Lab Immunol* 1996; **3**:590–3.
 39. Waltrip RW, Torrey EF, King LR, Weinberger DR, Rickler KC, Rubin SA, *et al.* Borna Disease Virus correlates in schizophrenia—monozygotic twin study cohort. *Soc Neurosci* 1995; **21**:748.
 40. Waltrip RW, Torrey EF, King LR, Weinberger DR, Rickler KC, Rubin SA, *et al.* Schizophrenia, Borna Disease, and neurological signs in a monozygotic twin study cohort. *Biolog Psych* 1996; **39**:500–666.
 41. Waltrip RW, Lieberman J, Robinson D, Bilder RM, Alvir JM, King LR, *et al.* First episode schizophrenia and Borna Disease Virus. *Schizophr Res* 1997; **24**:261.
 42. Waltrip RW, Lieberman J, Robinson D, Bilder RM, Alvir JM, King LR, *et al.* Borna Disease Virus seroconversion in first episode schizophrenia. *Biol Psychiatry* 1997; **41**:1–120S.
 43. Horimoto T, Takahashi H, Sakaguchi M, Horikoshi K, Iritani S, Kazamatsuri H, *et al.* A reverse-type sandwich enzyme-linked immunosorbent assay for detecting antibodies to Borna Disease Virus. *J of Clin Microbiol* 1997; **35**:1661–6.
 44. Nowotny N, Windhaber J. Borna Disease Virus and neuropsychiatric disorders. *Lancet* 1997; **350**:594.
 45. Chen C-H, Chiu Y-L, Wei F-C, Koong F-J, Liu H-C, Shaw C-K, *et al.* High seroprevalence of Borna virus infection in schizophrenic patients, family members and mental health workers in Taiwan. *Mol Psych* 1999; **4**:33–8.
 46. Deutz A, Fuchs K, Hinterdorfer F, Schuller W. Serological examination of veterinarians for zoonoses. Paper 1: Basic data and prevalence of antibodies against bacterial zoonoses. *Wien Tierarztl Mschr* 1996; **83**:283–8.
 47. Deutz A, Fuchs K, Nowotny N, Schuller W. Serological examination on zoonoses in veterinarians. Part III: Prevalence of antibodies to viral zoonoses and prophylaxis. *Wien Tierarztl Mschr* 1997; **84**:211–19.
 48. Nowotny N, Deutz A, Fuchs K, Schuller W, Hinterdorfer F, Auer H, *et al.* Prevalence of swine influenza and other viral, bacterial and parasitic zoonoses in veterinarians. *J Infect Dis* 1997; **176**:1414–15.
 49. Yamaguchi K, Sawada T, Tohru N, Igata-Yi R, Shiraki H, Horii Y, *et al.* Detection of Borna Disease virus-reactive antibodies from patients with psychiatric disorders and from horses by electrochemiluminescence immunoassay. *Clin Diag Lab Immunol* 1999; **6**:696–700.
 50. Nakaya T, Tada M, Takahashi H, Fujiwara S, Sakuma S, Sawamura Y, *et al.* Expression of Borna Disease Virus messages in clinical samples from patients with brain malignant tumours. *Proc. Japan Acad* 1996; **72(b)**:157–62.
 51. De la Torre JC, Gonzales-Dunia D, Cubitt B, Mallory M, Mueller-Lantzsch N, Grasser FA, *et al.* Detection of Borna Disease Virus antigen and RNA in human autopsy brain samples from neuropsychiatric patients. *Virology* 1996; **223**: 272–82.
 52. Salvatore M, Morzunov S, Schwemmler M, Lipkin W and the Borna Virus study Group. Borna disease virus in brains of North American and European people with schizophrenia and bipolar disorder. *Lancet* 1997; **349**:1813–14.
 53. Haga S, Motoi Y, Ikeda K, and the Japanese Bornavirus study group. Borna disease virus and neuropsychiatric disorders. *Lancet* 1997; **350**:593.
 54. Haga S, Yoshimura M, Motoi Y, Arima K, Aizawa T, Ikuta K, *et al.* Detection of Borna disease virus genome in normal human brain tissue. *Brain Res* 1997; **770**:307–9.
 55. Lieb K, Hallensleben W, Czygan M, Stitz L, Staeheli P, and the Bornavirus study Group. No Borna disease virus-specific RNA detected in blood from psychiatric patients in different regions of Germany. *Lancet* 1997; **350**:1002.
 56. Wolff C, Boomgaarden M, Koster-Eiserfunke W, Kleesiek K. Two years of experience with PCR screening of individual blood donations. *Infusion Therapy and Transfusion Medicine* 1998; **25**:128–34.
 57. Iwata Y, Takahashi K, Peng X, Fukuda K, Ohno K, Ogawa T *et al.*, Detection and sequence analysis of Borna disease virus p24 RNA from peripheral blood mononuclear cells of patients with mood disorders or schizophrenia and of blood donors. *J Virol* 1998; **72**:10044–9.
 58. Sierra-Honigmann AM, Carbone KM, Yolken RH. Polymerase chain reaction (PCR) search for viral nucleic acid sequences in schizophrenia. *Br J Psych* 1995; **166**:55–60.
 59. Kim YK, Kim SH, Choi SH, Ko YH, Kim L, Lee MS *et al.*, Failure to demonstrate Borna disease virus genome in peripheral blood mononuclear cells from psychiatric patients in Korea. *Journal of Neuro Virol* 1999; **5**:196–9.
 60. Shanker V, Dietzschold B, Fu ZF, Amsterdam J. A search for Bornavirus-specific genome in human nerve tissue by reverse transcriptase polymerase chain reaction (RT-PCR). *Biol Psych* 1992; **31**:103A.
 61. Bode L. Human infections with Borna Disease Virus and potential pathogenic implications. *Curr Top Microbiol Immunol* 1995; **190**:103–30.
 62. Bode L, Durrwald R, Rantam FA, Ferszt R, Ludwig H. First isolates of infectious human Borna Disease Virus from patients with mood disorders. *Mol Psych* 1996; **1**:200–12.

63. De la Torre JC, Bode L, Durrwald R, Cubitt B, Ludwig H. Sequence characterisation of human Borna Disease virus. *Virus Res* 1996; **44**:33–44.
64. Kishi M, Nakaya T, Yurie N, Kakinuma M, Takahashi TA, Sekiguchi S, et al. Prevalence of Borna disease virus RNA in peripheral blood mononuclear cells from blood donors. *Med Microbiol Immunol* 1995; **184**:135–8.
65. Kishi M, Nakaya T, Nakamura Y, Zhong Q, Ikeda K, Senjo M, et al. Demonstration of human Borna disease virus RNA in human peripheral blood mononuclear cells. *FEBS Letters* 1995; **364**:293–7.
66. Kishi M, Arimura Y, Ikuta K, Shoya Y, Lai PK, Kakinuma M. Sequence variability of Borna Disease Virus open reading frame II found in human peripheral blood mononuclear cells. *J Virol* 1996; **70**:635–40.
67. Kitani T, Kuratsune H, Fuke I, Nakamura Y, Nakaya T, Asahi S, et al. Possible correlation between Borna disease virus infection and Japanese patients with chronic fatigue syndrome. *Microbiol Immunol* 1996; **40**:459–62.
68. Nakaya T, Takahashi H, Nakamura Y, Asahi S, Tobiume M, Kuratsune H, et al. Demonstration of Borna disease virus RNA in peripheral blood mononuclear cells derived from Japanese patients with chronic fatigue syndrome. *FEMS Lett* 1996; **378**:145–9.
69. Igata-Yi R, Yamaguchi K, Yoshiki K, Takemoto S, Yamasaki H, Matsuoka M, et al. Borna disease virus and the consumption of raw horse meat. *Nat Med* 1996; **2**:948–9.
70. Sauder C, Muller A, Cubitt B, Mayer J, Steinmetz J, Trabert W, et al. Detection of Borna Disease Virus (BDV) antibodies and BDV RNA in psychiatric patients: evidence for high sequence conservation of human blood-derived BDV RNA. *J Virol* 1996; **70**:7713–24.
71. Kubo K, Fujiyoshi T, Yokoyama MM, Kamel K, Richt JA, Kitzte B, et al. Lack of association of Borna disease virus and human T-cell leukemia virus type 1 infections with psychiatric disorders among Japanese patients. *Clin Diag Lab Immunol* 1997; **4**:189–94.
72. Richt JA, Alexander RC, Herzog S, Hooper DC, Kean R, Spitsin S, et al. Failure to detect Borna disease virus infection in peripheral blood leukocytes from humans with psychiatric disorders. *J Neurol* 1997; **3**:174–8.
73. Gow JW, de la Torre JC, Behan WMH, Simpson K, McGill M, Dinan T, et al. Borna Disease virus in chronic fatigue syndrome. *Neurol Infect Epidemiol* 1997; **2**:63–6.
74. Gow J, Behan W, de la Torre J, Dinan T, McGill M, Simpson K, et al. Borna Disease virus and chronic fatigue syndrome. *Abstract in J Neurol Sci* 1997; **150**:S225.
75. Iwahashi K, Watanabe M, Nakamura K, Fujiwara Y, Isojima G, Suwaki H, et al. The relationship between Borna Disease Virus Infection and Schizophrenia. *Clin Psych* 1997; **39**: 1275–9.
76. Iwahashi K, Watanabe M, Nakamura K, Suwaki H, Nakaya T, Nakamura Y, et al. Clinical investigation of the relationship between Borna disease virus (BDV) infection and schizophrenia in 67 patients in Japan. *Acta Psychiatrica Scand* 1997; **96**:412–15.
77. Iwahashi K, Watanabe M, Nakamura K, Suwaki H, Nakaya T, Nakamura Y, et al. Positive and negative syndromes, and Borna disease virus infection in schizophrenia. *Neuropsychobiol* 1998; **37**:59–64.
78. Iwahashi K, Watanabe M, Nakamura K, Suwaki H, Nakaya T, Nakamura Y, et al. Borna disease virus infection and negative syndromes in Japanese schizophrenia patients. *Psych Clin Neurosci* 1998; **32**:119.
79. Deuschle M, Heuser I, Gotthardt U, Schmider J, Bode L, Ludwig H. Borna Disease virus antigen in cerebrospinal fluid of patients with recurrent depression and multiple sclerosis. *Pharmacopsychiat* 1997; **30**:169 (abstracts of the 20th symposium of AGNP, Nuremberg 1997).
80. Deuschle M, Bode L, Heuser I, Schmider J, Ludwig H. Borna disease virus proteins in cerebrospinal fluid of patients with recurrent depression and multiple sclerosis. *Lancet* 1998; **352**:1828–9.
81. Igata T, Yamaguchi K, Igata-Yi R, Yoshiki K, Takemoto S, Yamasaki H, et al. Dementia and Borna disease virus. *Dement Geriatric Cognitive Dis* 1997; **9**:24–5.
82. Seltén JP, van Loon AM, van Vliet K, Pleyte W, Hoek HW, Kahn RS. Borna Disease Virus in Caribbean immigrants to the Netherlands, diagnosed with schizophrenia. *Schizophrenia Res* 1998; **29**:1–2 (abstract).
83. Takahashi H, Nakaya T, Nakamura Y, Asahi S, Onishi Y, Ikebuchi K, et al. Higher prevalence of Borna disease virus infection in blood donors living near thoroughbred horse farms. *J Med Virol* 1997; **52**:330–5.
84. Bechter K, Schüttler R, Herzog S. Case of Neurological and behavioral abnormalities: due to Borna Disease Virus Encephalitis. *Psych Res* 1992; **42**:193–6.
85. Bechter K, Schüttler R, Herzog. Borna Disease Virus: possible causal agent in Psychiatric and neurological disorders in two families. *Psych Res* 1992; **42**:291–4.
86. Nowotny N, Dürr G. Enteroviral hypothesis for motor neurone disease. *Br Med J* 1995; **310**:256.
87. Planz O, Rentzch C, Batra A, Rziha H-J, Stitz L. Persistence of Borna Disease virus-specific nucleic acid in blood of psychiatric patients. *Lancet* 1998; **352**:623.
88. Rothman KJ & Greenland S. Causation and Causal Inference. In: Rothman KJ, Greenland S, eds. *Modern Epidemiology*, 2nd edn. Philadelphia, Lipincott Williams & Wilkins, 1998.
89. Deeks JJ. Systematic reviews of published evidence: Miracles or minefields? *Ann Oncol* 1998; **9**:1–7.
90. Schwemmler M, Jehle C, Formella S, Staeheli P. Sequence similarities between human bornavirus isolates and laboratory strains question human origin. *Lancet* 1999; **354**: 1973–4.
91. Bode L, Stoyloff R, Ludwig H. Human bornaviruses and laboratory strains. *Lancet* 2000; **355**:1462.
92. Nowotny N, Kolodziejek. Demonstration of Borna Disease Virus Nucleic Acid in a patient with Chronic Fatigue Syndrome. *J Infect Dis* 2000; **181**:1860–1.